

Erratum: This table replaces Table 1 on p. 596. Changes are in boldface type.

Table 1. Weighted mean (standard error) density of snags by wildlife habitat type, successional stage, and snag size, Oregon and Washington.¹

Wildlife habitat type	<i>Successional Stage and Snag Size</i>							
	<i>Early</i>		<i>Middle</i>		<i>Late</i>		<i>All stages</i>	
	<i>Total</i>	<i>Large</i>	<i>Total</i>	<i>Large</i>	<i>Total</i>	<i>Large</i>	<i>Total</i>	<i>Large</i>
	<i>Mean (SE) trees per acre</i>							
Westside conifer-hardwood	2.1 ^a (0.1)	0.8 ^a (0.0)	7.2^b (0.2)	2.4^b (0.1)	12.7^c (0.4)	5.8^c (0.2)	5.8 (0.1)	2.2 (0.1)
Westside white oak-Douglas-fir	2.5 ^a (0.4)	0.6 ^a (0.2)	4.6^a (0.4)	1.1^a (0.1)	6.9^b (1.0)	2.4^b (0.4)	4.1 (0.3)	1.0 (0.1)
SW OR mixed conifer-hardwood	3.8 ^a (0.4)	1.0 ^a (0.1)	6.9^b (0.4)	2.0^b (0.1)	8.5^b (0.4)	3.8^c (0.2)	6.2 (0.2)	2.1 (0.1)
Montane mixed-conifer	7.2 ^a (0.6)	1.2 ^a (0.1)	20.0^b (0.5)	4.2^b (0.2)	16.3^c (0.6)	8.8^c (0.3)	15.1 (0.4)	3.9 (0.1)
Subalpine parkland	14.1 (3.0)	0.7 ^a (0.2)	15.1 (1.7)	2.3^b (0.4)	NA	NA	14.6 (1.7)	1.5 (0.2)
Eastside mixed-conifer	6.0 ^a (0.4)	0.8 ^a (0.1)	8.7 ^b (0.2)	1.7 ^b (0.0)	8.4 (0.2)	3.2 ^c (0.2)	7.9 (0.2)	1.5 (0.0)
Lodgepole pine	6.7 ^a (0.6)	0.3 ^a (0.0)	11.2 ^b (1.1)	0.9 ^b (0.1)	NA	NA	8.0 (0.5)	0.5 (0.0)
Ponderosa pine (eastside)	2.0 (0.2)	0.4 (0.0)	2.0 (0.2)	1.1 (0.0)	2.1 (0.3)	0.6 (0.1)	2.0 (0.1)	0.4 (0.0)
Western juniper	0.3 (0.1)	0.1 (0.0)	0.6 (0.2)	0.2 (0.1)	0.3 (0.2)	0.1 (0.1)	0.3 (0.1)	0.1 (0.0)
All wildlife habitat types	3.7 ^a (0.1)	0.8 ^a (0.0)	8.6 ^b (0.1)	2.1 ^b (0.0)	11.5 ^c (0.2)	5.4 ^c (0.1)	7.0 (0.1)	2.0 (0.0)

¹ NA: Not applicable -- sample size <10 plots. Significantly different means (alpha=0.05) within rows (among successional stages for a given snag size-class) are indicated by different letter footnotes. "Total" includes snags >= 10.0 in DBH, decay classes 1-5, and >=6.6 ft tall; "large" includes snags >=19.7 in DBH, decay classes 1-5, and >=6.6 ft tall. Table modified from Ohmann and Waddell (in press).

Erratum: This table replaces Table 2 on p. 597. Changes are in boldface type.

Table 2. Weighted mean (standard error) volume, percent cover, and density of total down wood by wildlife habitat type and successional stage, Oregon and Washington.¹

Wildlife habitat type	Successional Stage											
	Early	Middle	Late	All stages	Early	Middle	Late	All stages	Early	Middle	Late	All stages
	<i>Mean (SE) cubic feet per acre</i>				<i>Mean (SE) percent cover</i>				<i>Mean (SE) pieces per acre</i>			
Westside conifer-hardwood	2,169 ^a (109)	2,396 ^a (89)	3,234 ^b (120)	2,619 (61)	4.4 ^a (0.2)	4.5 ^a (0.1)	5.5 ^b (0.2)	4.8 (0.1)	132.1 ^a (5.6)	102.9^a (2.8)	101.5 ^b (2.8)	109.7 (2.1)
Westside white oak-Douglas-fir	772 (167)	632 (87)	1,017 (176)	733 (73)	1.8 (0.3)	1.5 (0.1)	2.2 (0.3)	1.7 (0.1)	70.5 (11.1)	41.1 (4.1)	51.9 (7.5)	48.9 (3.8)
SW OR mixed conifer-hardwood	1,213 ^a (123)	983 ^a (66)	1,695 ^b (114)	1,227 (54)	2.5 ^a (0.2)	2.2 ^a (0.1)	3.2 ^b (0.2)	2.5 (0.1)	64.3 ^a (5.9)	59.7^a (3.2)	69.7 ^b (3.8)	63.5 (2.3)
Montane mixed-conifer	1,459 ^a (91)	1,608 ^a (47)	2,837 ^b (151)	1,769 (46)	3.8 ^a (0.2)	4.3 ^b (0.1)	5.1 ^c (0.2)	4.3 (0.1)	102.8 ^a (4.1)	101.3^b (2.6)	90.0 ^c (3.8)	99.8 (1.9)
Subalpine parkland	502 (104)	789 (149)	NA	629 (90)	1.6 (0.3)	2.1 (0.4)	NA	1.8 (0.2)	45.8 (7.0)	52.2 (7.9)	NA	47.8 (5.1)
Eastside mixed-conifer	672 ^a (31)	780 ^b (19)	840 ^b (70)	753 (16)	2.0 (0.1)	2.2 ^a (0.0)	1.9 ^b (0.1)	2.1 (0.0)	63.9 (2.0)	57.8^a (1.1)	41.3 ^b (2.7)	58.5 (0.9)
Lodgepole pine	714 (36)	792 (60)	NA	734 (31)	2.9 (0.1)	2.8 (0.2)	NA	2.9 (0.1)	81.6 (3.4)	65.2 (5.0)	NA	77.3 (2.8)
Ponderosa pine (eastside)	304 ^a (19)	406 ^b (20)	282 (43)	362 (14)	0.8 ^a (0.0)	1.0 ^b (0.0)	0.7 ^a (0.1)	0.9 (0.0)	28.4 ^a (1.6)	31.9^b (1.2)	18.1 ^a (2.8)	29.7 (0.9)
Western juniper	107 (44)	104 (27)	107 (67)	106 (27)	0.2 (0.1)	0.3 (0.1)	0.3 (0.2)	0.2 (0.0)	6.6 (1.7)	9.3 (2.5)	10.9 (6.6)	7.8 (1.4)
All wildlife habitat types	1,000 ^a (27)	1,109 ^b (19)	2,306^c (64)	1,255 (17)	2.6 ^a (0.1)	2.7 ^b (0.0)	4.1 ^c (0.1)	2.9 (0.0)	74.0 ^a (1.4)	67.7^b (0.8)	77.4^c (1.6)	71.0 (0.7)

¹ NA: Not applicable -- sample size <10 plots. Total down wood includes pieces ≥ 4.9 in large end diameter, decay classes 1-4, and ≥ 6.6 ft long. Significantly different means (alpha = 0.05) among successional stages are indicated by different letter footnotes. Table modified from Ohmann and Waddell (in press).

Erratum: This table replaces Table 3 on p. 598. Changes are in boldface type.

Table 3. Weighted mean (standard error) volume, percent cover, and density of large down wood by wildlife habitat type and successional stage, Oregon and Washington.¹

Wildlife habitat type	Successional Stage				Successional Stage				Successional Stage			
	Early	Middle	Late	All stages	Early	Middle	Late	All stages	Early	Middle	Late	All stages
	<i>Mean (SE) cubic feet per acre</i>				<i>Mean (SE) percent cover</i>				<i>Mean (SE) pieces per acre</i>			
Westside conifer-hardwood	1,408 ^a (97)	1,709 ^a (81)	2,459^b (114)	1,883 (57)	1.9 ^a (0.1)	2.2 ^a (0.1)	3.1 ^b (0.1)	2.4 (0.1)	23.9 ^a (1.6)	19.7 ^a (0.9)	24.1 ^b (1.1)	22.3 (0.7)
Westside white oak-Douglas-fir	442 (147)	373 (79)	609 (154)	432 (64)	0.6 (0.2)	0.6 (0.1)	0.9 (0.2)	0.6 (0.1)	9.0 (2.4)	5.9 (1.1)	6.5 (1.8)	6.6 (0.9)
SW OR mixed conifer-hardwood	766 ^a (107)	609 ^a (57)	1,197 ^b (106)	803 (49)	0.9 ^a (0.1)	0.9 ^a (0.1)	1.6 ^b (0.1)	1.1 (0.1)	7.9 ^a (1.2)	7.4 ^a (0.8)	12.0 ^b (1.2)	8.9 (0.6)
Montane mixed-conifer	699 ^a (80)	660 ^a (34)	2,071 ^b (137)	906 (40)	1.0 ^a (0.1)	1.0 ^a (0.0)	2.9 ^b (0.2)	1.3 (0.1)	10.3 ^a (0.9)	8.7 ^a (0.4)	23.4 ^b (1.5)	11.6 (0.4)
Subalpine parkland	104 (47)	319 (91)	NA	207 (54)	0.2 (0.1)	0.5 (0.1)	NA	0.4 (0.1)	2.6 (1.2)	4.4 (1.4)	NA	3.4 (0.9)
Eastside mixed-conifer	247 ^a (26)	329 ^b (14)	513 ^c (60)	317 (13)	0.4 ^a (0.0)	0.5 ^b (0.0)	0.7 ^c (0.1)	0.5 (0.0)	3.6 ^a (0.2)	4.2 ^b (0.2)	6.4 ^c (0.7)	4.2 (0.1)
Lodgepole pine	80 ^a (13)	209 ^b (40)	NA	116 (14)	0.2 ^a (0.0)	0.3 ^b (0.1)	NA	0.2 (0.0)	1.7 ^a (0.3)	3.2 ^b (0.7)	NA	2.1 (0.3)
Ponderosa pine (eastside)	147 ^a (16)	221 ^b (17)	160 (34)	191 (11)	0.2 ^a (0.0)	0.3 ^b (0.0)	0.3 (0.1)	0.3 (0.0)	2.3 ^a (0.2)	2.9 ^b (0.2)	2.5 (0.6)	2.7 (0.2)
Western juniper	71 (40)	50 (23)	89 (66)	64 (24)	0.1 (0.0)	0.1 (0.0)	0.2 (0.2)	0.1 (0.0)	0.4 (0.2)	0.8 (0.4)	4.5 (3.3)	0.7 (0.2)
All wildlife habitat types	496 ^a (23)	579 ^b (16)	1,695 ^c (59)	720 (14)	0.7 ^a (0.0)	0.8 ^b (0.0)	2.2 ^c (0.1)	1.0 (0.0)	7.1 ^a (0.3)	6.9 ^b (0.2)	17.5 ^c (0.6)	8.5 (0.2)

¹ NA: Not applicable -- sample size <10 plots. Large down wood includes pieces ≥ 19.7 in large end diameter, decay classes 1-4, and ≥ 6.6 ft long. Significantly different means ($\alpha = 0.05$) among successional stages are indicated by different letter footnotes. Table modified from Ohmann and Waddell (in press).